



Chinese Privet Control with Herbicide Foliar Sprays

by James H. Miller, USDA Forest Service, Southern Research Station, Auburn, AL 36849

Chinese privet (*Ligustrum sinense* Lour.) is probably the most problematic alien shrub in the 13-states of the southern region, widely invading forests, parks and preserves, pastures, and right-of-ways (Haragan 1996, Miller 1997, Matlack 2002). After a century of planting as an ornamental shrub following introduction in 1852 (Dirr 1998), range expansion has been rapid and far-reaching since about 1960 (NRCS website). This rapid invasion has occurred as birds feed upon abundant fruits produced in early spring and disperse seed during northern migrations. It is widely observed that the habitats most under siege are disturbed areas and bottomland forests (Dirr 1998), while upland forests and pasture margins are steadily being invaded as well. Site dominance occurs through Chinese privet's production of abundant root suckers and clump sprouts, as well as carpets of seedlings in infested areas.

Chinese privet is but one of at least eight nonnative privet species within the region that have escaped into natural habitats. Other widely occurring privet invaders are European privet (*L. vulgare* L.), glossy privet (*L. lucidum* Ait. f.), and Japanese privet (*L. japonicum* Thunb.). Those locally problematic are Amur privet (*L. amuense* Carr.), border privet (*L. obtusifolium* Sieb. & Zucc.), California (originally from Japan) privet (*L. ovalifolium* Hassk.), and waxyleaf privet (*L. quihoui* Carr.), including several varieties of each. Japanese, glossy, and border privet are evergreen while the others are semi-evergreen and retain foliage depending on the severity of the winter and locale. All have opposite leaves,

with white flower clusters in spring that yield black to blue-black drupes in fall and winter to spring. The fleshy one-seeded fruit characterize these members of the olive family (Oleaceae).

Herbicides are one tool that can be used to control privets as part of an integrated vegetation management approach. The objective of this investigation was to compare foliar sprays of most herbicides registered for forest use in the southern region for their effectiveness on Chinese privet.

Methods

The study site was located along a riparian area of a perennial stream in east-central Alabama. A uniformly dense stand of Chinese privet had been brush mowed on a 3-year cycle for 9 years before study initiation. The infestation had one year of regrowth, being 4 to 10 ft tall. Sixty-four, 10 x 20-ft plots were established in four blocks. Seven herbicide treatments and a non-treated control were randomly assigned to plots in each block and tested at two intervals, August and September. The first treatment was planned for July, but due to the absence of rainfall for 3 months during the summer, it was applied in August after rainfall commenced with 2 inches in 2 weeks preceding treatment.

Near maximum labeled rates were tested for each herbicide using formulations with a single active ingredient (Table 1). This approach aimed to identify the most effective active ingredients for treating Chinese privet in late-summer and early fall. Applications were by a CO₂-powered backpack sprayer with a

Table 1. Herbicide tests on Chinese privet.

Herbicide active ingredient (ai)	Rate ¹ per Acre lbs ai ²	1 Year After Treatment	2 Years After Treatment	3 Years After Treatment
Accord glyphosate	1.5 gal 6.0	99a ³	98a	97a
Arsenal AC imazapyr	24 fl oz 0.75	94a	89a	79ab
Escort metsulfuron	3.3 oz 0.12	81ab	79a	69ab
Garlon 4 triclopyr	1.5 gal 6.0	64ab	44ab	22abc
Oust sulfometuron	6.0 oz 0.28	31abc	32abc	21abc
Vanquish dicamba	1.5 gal 6.0	27abc	25abc	04abcd
Tordon K picloram	0.5 gal 1.0	12abcd	09abcd	05abcd
Transline clopyralid	21 fl oz 0.5	00abcd	00abcd	00abcd

¹Product per acre. ²Pounds active ingredient per acre. ³Results of Tukey's HSD, where values with different letters are significantly different at the 5% level of probability.



Spraying Systems XR 8003 flat-fan nozzle swiveled downward on a 4-foot extended wand. A total spray mixture of 40 gallons per acre (gpa) in water was used except with Accord, which was tested with 30 gpa per label recommendations. A surfactant (Entry II) was added at 0.5 percent to all mixtures except Accord, which contains surfactant. The applicator stood in the middle of half plots and rotated about to uniformly cover all plants, while a guide outside the plot gave directions to assure uniformity (shown above).

Plots were rated 1, 2, and 3 years after treatment using visual estimates of percent volume reduction of standing shrubs. Untreated control plots in each block were used as height references during ratings, while before treatment volumes (average height X cover) were used as co-variants in the analysis. There were no significant differences between


the August and September applications and thus these data were combined. According to standard procedures, percent reductions (control) were arsine square root transformed and data were analyzed using Tukey's HSD Test.

Findings

Privet control exceeded 90 percent with Accord (glyphosate) and Arsenal AC (imazapyr) the first year and did not statistically differ from Escort, which averaged 81 percent control (Table 1). Accord gave near complete control for the 3-year post-treatment period to exceed 97 percent with minimal resprouting and seedlings. Garlon 4, Oust, Vanquish, Tordon K, and Transline provided less than 65 percent control. Transline was completely ineffective on privet at these timings and rates.

The most effective herbicide, Accord, is a foliar active herbicide that is deactivated when it reaches the soil and thus presents safety to nearby unsprayed plants. This offers a treatment option to prevent harm to native cohort plants when care in application restricts spray to privet foliage only. The wand extension used in this research permitted effective treatment for privet that was 10 ft tall. The wand could be fitted with a longer extension and a projecting spray tip to treat taller privet.

The active ingredient in Accord is glyphosate, which is available in many formulations including aquatic labeled products permitted for spraying around and over water. The aquatic formulations could be used to eradicate the extensive privet infestations that occur along streamside areas and



partially flooded wetlands. Further tests of aquatic formulations of glyphosate are required since they contain no surfactants, which may lessen control.

Conclusions

Resprouted Chinese privet can be effectively controlled or even eradicated with Accord treatments or other herbicides with similar glyphosate formulations. This offers a treatment option that can be safe to applicators as well as flora and fauna when used according to label directions. Subsequent tests have shown that lower rates are equally effective, as low as 1 quart per acre. Treatments in December are most effective while those in April are only slightly less effective (Harrington and Miller 2005). Arsenal AC and Escort were less effective in this trial, but provided enough control for use in particular locations. All treatments will need to be repeated to achieve eradication, and native plant revegetation fostered to obtain restoration.

Literature

Dirr, M.A. 1998. Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation, and uses. Stipes Publication, Champaign, IL. 1187 p.

Haragran, D.P. 1996. *Ligustrum vulgare*, *L. sinense*, *L. japonicum*. p 58-59. In: Randall, J.M and Marinelli, J (Eds.) Invasive plants: weeds of the global garden. Brooklyn Botanic Garden, NYC. 111 p.

Harrington, T. and Miller, J.H. 2005. Effects of application rate, timing, and formulation of glyphosate and triclopyr for control of Chinese privet (*Ligustrum sinense*). Weed Technology 19:47-54.

Matlack, G.R. 2002. Exotic plant species in Mississippi, USA: critical issues in management and research. Natural Areas Journal 22:241-247.

Miller, J.H. 1997. Exotic invasive plants in southeastern forests. p 97-105. In: K.O. Britton (Ed.), Proceedings of Exotic pests of eastern forests. USDA Forest Service and Tennessee Exotic Pest Plant Council. 198 p.



Intelligent Solutions.

Dedicated Service.




Garlon* 3A specialty herbicide • Garlon* 4 specialty herbicide
Rodeo® herbicide • DMA* 4 IVM herbicide

- Proven performance
- Returnable, refillable containers
- Experienced sales force

For more information, contact Scott Ditmarsen
(813-866-7090 or scditmarsen@dow.com)
or visit

www.vegetationmgmt.com

*Trademark of Dow AgroSciences LLC
Always read and follow label directions



Solutions for Invasive Weed Control

Officers

Jim Burney, Chair
Aquatic Vegetation Control, Inc.
561/845-5525
L1J2@aol.com

Dianne Owen, Secretary
Florida Atlantic University
954-236-1085
dowen@fau.edu

Kristina Kay Serbesoff-King,
Treasurer
The Nature Conservancy
561-744-6668
kserbesoffking@tnc.org

Karen Brown, Editor
University of Florida
Center for Aquatic
& Invasive Plants
352/392-1799
kpbrown@mail.ifas.ufl.edu

Mike Bodle,
Immediate Past Chair
South Florida Water
Management District
561/682-6132
mbodle@sfwmd.gov

Alison Fox, Chair Elect
University of Florida
Agronomy Department
352/392-1811, Ext. 207
amfox@ifas.ufl.edu

Directors

Roger Clark (2nd year)
Lee County Parks & Recreation
239/461-7453
Roger@leegov.com

Drew Leslie (2nd year)
Florida DEP Bureau of Invasive
Plant Management
850/245-2822
Drew.Leslie@dep.state.fl.us

Cressida Silvers (2nd year)
USDA/ARS
954/475-0541 ext. 144
csilvers@saa.ars.usda.gov

Jim Duquesnel (2nd year)
Florida Park Service
305/451-1226
james.g.duquesnel@dep.state.fl.us

Scott Ditmarsen (1st year)
DowAgro Sciences
813/866-7090
scditmarsen@dow.com

Jon Lane (1st year)
USACOE Invasive Species
Management
904/232-1044
Jon.S.Lane@saj02.usace.army.mil

Tony Pernas (1st year)
National Park Service
Florida/Caribbean Exotic Plant
Management Team
305/252-0347
Tony_Pernas@nps.gov

Paul Pratt (1st year)
USDA-ARS Invasive Plant
Research Laboratory
954/475-6549
prattp@saa.ars.usda.gov

Committee Chairs

By-laws
Dennis Giardina
941/657-7637
Dennis_Giardina@fws.gov

CAST Representative
Dennis Giardina
Ken Langeland (alternate)

Editorial
Karen Brown

Education
Leesa Souto
Stormwater Management
Academy
321/722-2123
Lsouto@mail.ucf.edu

FNGLA/FLEPPC Liaison
Doria Gordon
University of Florida
The Nature Conservancy
352/392-5949
dgordon@botanyufl.edu
-and-
JB Miller
St. Johns River Water
Management District
millerjb@aug.com

Legislative
Matthew King
Palm Beach County
561/233-2400
mking@co.palm-beach.fl.us

Local Arrangements
Mike Bodle

Membership
Katy Roberts
727/726-1455
kroberts@j.net

Merchandise
Tony Pernas

Nominations
Mike Bodle

Outreach
Tom Fucigna
CZR Inc.
561/747-7455
tomfucignajr@hotmail.com

Plant List
Kathy Craddock Burks
Florida Natural Areas
Inventory
850/224-8207 Ext. 210
kburks@fna.org

Program Chair
Daniel Clark
Florida/Caribbean Exotic Plant
Management Team
340/693-8950 x 231
daniel_clark@nps.gov

Research
John Volin
Florida Atlantic University
954/236-1115
jvolin@fau.edu

Training
Jim Duquesnel

Vendors
Scott Ditmarsen

Webmaster
Tony Pernas

Task Force Chairs

Australian Pine
Robert Egan
Habitat Restoration Resources
239/574-8173
HabitatRR@yahoo.com

Brazilian Pepper
Jim Cuda
University of Florida
Entomology Department
352/392-1901 Ext. 126
Jcuda@ifas.ufl.edu
Carrotwood
Chris Lockhart
Habitat Specialists, Inc.
561/738-1179
chris@habitat specialists.com

Dioscorea
Mike Bodle

Grasses
Greg MacDonald
University of Florida
Agronomy Department
352/392-1811 Ext. 228
gemac@mail.ifas.ufl.edu

Lygodium
Amy Ferriter/Tom Fucigna
Shunkvine
Brian Nelson
SWFWMD
352/796-7211
Brian.Nelson@swfwmd.state.fl.us

Chinese Tallow
Drew Leslie

Melaleuca
Francois Laroche
South Florida Water
Management District
561/682-6193
flaroche@sfwmd.gov

SOUTHEAST EXOTIC PEST PLANT COUNCIL www.se-eppc.org

Officers

President
Brian Bowen
Tennessee Dept Environment
and Conservation
Division of Natural Heritage
615/532-0436
Brian.Bowen@state.tn.us

Vice President
Joyce Bender
Kentucky State Nature
Preserves Commission
502/573-2886
Joyce.Bender@ky.us

Secretary
Kristen Gounaris Allen
Richmond National
Battlefield Park
804/795-5019
Kristen_Allen@nps.gov

Treasurer
Tony Pernas, Florida EPPC

**SE-EPPC Chapters and
Regional Organizations:**
Alabama, Florida, Georgia,
Kentucky, Mid-Atlantic,
Mississippi, New England,
North Carolina, South Carolina,
Tennessee, USDA-Forest Service

Wildland Weeds

SUMMER 2005, VOLUME 8, NUMBER 3

Table of Contents

- 4 Editor's Note
- 5 Chinese Privet Control with Herbicide Foliar Sprays
by James H. Miller
- 9 Melapaleuza 2005 Rocked! by Cressida Silvers
- 11 Invasive Plant Lists and Assessments – A Brief Introduction
- 13 FLEPPC 2005 List of Invasive Species
- 17 Putting a Stop to the Cat-claw Vine Infestation in
Gainesville by Daniel B. Ward
- 18 A New Approach to Combat Invasive Species
by Susan Musante
- 20 Palm Beach County Ordinance Helps Residents Fight
Invasive Plants by Matthew King
- 22 Just Who is Daniel W. Clark?
- 23 A Farewell to Ferriter
- 24 New Quarantine Facility Takes Flight by Cressida Silvers
- 25 Headway on Capitol Hill Against Invasive Species
- 27 Internodes

The mission of the Florida Exotic Pest Plant Council is to support the management of invasive exotic plants in Florida's natural areas by providing a forum for the exchange of scientific, educational and technical information.

An exotic plant has been introduced to Florida, either purposefully or accidentally, from a natural range outside of Florida. A **naturalized exotic plant** is one that sustains itself outside of cultivation (it is still exotic; it has not "become" native). An **invasive exotic plant** not only has become naturalized, but it is expanding its range in Florida plant communities.

Wildland Weeds (ISSN 1524-9786) is published quarterly by the Florida Exotic Pest Plant Council (FLEPPC) and the Southeast Exotic Pest Plant Council (SE-EPPC) to provide a focus for the issues and for information on exotic pest plant biology, distribution and control.

Direct all editorial and advertising inquiries to:

Karen Brown, Editor
Wildland Weeds
7922 NW 71st Street
Gainesville, FL 32653
352/392-1799; FAX 352/392-3462
kpbrown@ifas.ufl.edu

Direct address changes to:

Jackie Smith
DEP - Invasive Plant Management
9737 Gumbo Limbo Lane
Jensen Beach, FL 34957
561/722-2479
jackie.c.smith@dep.state.fl.us



Editorial Committee:

Mike Bodle
Kathy Craddock Burks
Jim Cuda
Tom Fucigna
Ken Langeland
Michael Meisenburg

Design by JS Design Studio.
Printing by StorterChilds,
Gainesville, FL.

The Exotic Pest Plant Council has not tested any of the products advertised or referred to in this publication, nor has it verified any of the statements made in any of the advertisements or articles. The Council does not warrant, expressly or implied, the fitness of any product advertised or the suitability of any advice or statements contained herein.

On the Cover: Melapaleuza turned into Mudapaleuza when 3 inches of rain fell two days prior to the event in the Everglades buffer strip near Holiday Park. Five swamp buggies got bogged down in the mud. See article on page 9. (Photo by Ken Langeland)